| Microorganism | 2013 | 2014 | 2015 | 2016 | 2017 | 2018 | 2019 | Total |
|--------------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-----------------|-------------------|
| S. arlettae | - | - | 1 (0.0%) | - | - | - | - | 1 (0.0%) |
| S. auricularis | 3 (0.2%) | 8 (0.3%) | - | 3 (0.1%) | 7 (0.2%) | 3 (0.1%) | 6 (0.2%) | 30 (0.2%) |
| S. capitis | 163 (8.2%) | 211 (8.0%) | 259 (9.1%) | 235 (7.8%) | 240 (8.1%) | 262 (8.2%) | 276 (8.8%) | 1,646 (8.3%) |
| S. caprae | 6 (0.3%) | 5 (0.2%) | 4 (0.1%) | 16 (0.5%) | 1 (0.0%) | 11 (0.3%) | 6 (0.2%) | 49 (0.2%) |
| S. carnosus | 1 (0.1%) | = | 2 (0.1%) | = | - | - | = | 3 (0.0%) |
| S. chromogenes | - | 1 (0.0%) | = | = | 2 (0.1%) | - | = | 3 (0.0%) |
| S. cohnii | 9 (0.5%) | 6 (0.2%) | 7 (0.2%) | 11 (0.4%) | 5 (0.2%) | 5 (0.2%) | 4 (0.1%) | 47 (0.2%) |
| S. condimenti | = | = | = | = | - | 2 (0.1%) | 1 (0.0%) | 3 (0.0%) |
| S. epidermidis | 1,024 (51.5%) | 1,365 (51.7%) | 1,557 (54.5%) | 1,546 (51.2%) | 1,544 (51.9%) | 1,606 (50.3%) | 1,543 (49.2%) | 10,185 (51.4%) |
| S. equorum | - | - | = | 1 (0.0%) | 3 (0.1%) | - | 1 (0.0%) | 5 (0.0%) |
| S. gallinarum | - | - | - | 1 (0.0%) | - | - | - | 1 (0.0%) |
| S. haemolyticus | 72(3.6%) | 130 (4.9%) | 141 (4.9%) | 154 (5.1%) | 145 (4.9%) | 198 (6.2%) | 141 (4.5%) | 981 (5.0%) |
| S. hominis | 631 (31.8%) | 816 (30.9%) | 789 (27.6%) | 945 (31.3%) | 934 (31.4%) | 1,009 (31.6%) | 1,037 (33.1%) | 6,161 (31.1%) |
| S. kloosii | - | - | - | 1 (0.0%) | - | - | - | 1 (0.0%) |
| S. lentus | - | = | = | 1 (0.0%) | - | 1 (0.0%) | 1 (0.0%) | 3 (0.0%) |
| S. lugdunensis | 27 (1.4%) | 25 (0.9%) | 11 (0.4%) | 19 (0.6%) | 17 (0.6%) | 23 (0.7%) | 31 (1.0%) | 153 (0.8%) |
| S. massiliensis | - | 4 (0.2%) | - | - | - | - | - | 4 (0.0%) |
| S. pasteuri | 3 (0.2%) | 4 (0.2%) | 6 (0.2%) | 3 (0.1%) | 3 (0.1%) | 4 (0.1%) | 9 (0.3%) | 32 (0.2%) |
| S. pettenkoferi | 3 (0.2%) | 2 (0.1%) | 3 (0.1%) | 2 (0.1%) | 2 (0.1%) | 13 (0.4%) | 22 (0.7%) | 47 (0.2%) |
| S. piscifermentans | - | - | - | - | - | 1 (0.0%) | 2 (0.1%) | 3 (0.0%) |
| S. saccharolyticus | = | 1 (0.0%) | 4 (0.1%) | 1 (0.0%) | 5 (0.2%) | 2 (0.1%) | 5 (0.2%) | 18 (0.1%) |
| S. saprophyticus | 2 (0.1%) | 12 (0.5%) | 7 (0.2%) | 16 (0.5%) | 10 (0.3%) | 6 (0.2%) | 3 (0.1%) | 56 (0.3%) |
| S. schleiferi | 7 (0.4%) | 6 (0.2%) | 17 (0.6%) | 7 (0.2%) | 4 (0.1%) | 3 (0.1%) | 4 (0.1%) | 48 (0.2%) |
| S. sciuri | - | 1 (0.0%) | 1 (0.0%) | 2 (0.1%) | - | - | - | 4 (0.0%) |
| S. simulans | 4 (0.2%) | 1 (0.0%) | 3 (0.1%) | 4 (0.1%) | 4 (0.1%) | 6 (0.2%) | 6 (0.2%) | 28 (0.1%) |
| S. warneri | 31 (1.6%) | 39 (1.5%) | 42 (1.5%) | 49 (1.6%) | 46 (1.5%) | 38 (1.2%) | 38 (1.2%) | 283 (1.4%) |
| S. xylosus | 1 (0.1%) | 2 (0.1%) | 1 (0.0%) | = | 3 (0.1%) | - | 1 (0.0%) | 8 (0.0%) |
| Total | 1,987 (100%) | 2,639 (100%) | 2,855 (100%) | 3,017 (100%) | 2,975 (100%) | 3,193 (100%) | 3,137 (100%) | 19,803 (100%) |