

| Antibiotic class           | Antibiotic                                 | S. caprae         | S. colinii          | S. epidemidis       | S. haemolyticus | S. hominis          | S. lugdunensis      | S. pettenkoferi | S. saprophyticus | S. schleiferi | S. warneri      |
|----------------------------|--|-------------------|---------------------|---------------------|-----------------|---------------------|---------------------|-----------------|------------------|---------------|-----------------|
| Aminoglycosides            | Gentamicin (GEN, J01GB03)                  | 2.6% (1/39)       | 0.0% (0/38)         | 40.7% (2,744/6,739) | 62.7% (352/561) | 7.3% (333/4,553)    | 3.7% (3/81)         | 0.0% (0/37)     | 0.0% (0/39)      | 0.0% (0/34)   | 13.2% (30/228)  |
|                            | Kanamycin (KAN, J01GB04)                   | 34.3% (79/230)    | 80.3% (3,174/3,954) | 80.3% (3,174/3,954) | 89.7% (358/399) | 67.0% (819/1,222)   | 18.1% (816/4,518)   | 3.7% (3/82)     | 0.0% (0/38)      | 0.0% (0/34)   | 41.9% (31/74)   |
|                            | Tobramycin (TOB, J01GB01)                  | 6.6% (79/1,198)   | 2.9% (1/35)         | 47.4% (3,167/6,686) | 63.1% (355/563) | 18.1% (816/4,518)   | 3.7% (3/82)         | 0.0% (0/35)     | 0.0% (0/38)      | 0.0% (0/34)   | 13.0% (30/230)  |
|                            | Chloramphenicol (CHL, J01BA01)             | 0.4% (3/817)      | 1.6% (75/4,662)     | 1.6% (75/4,662)     | 4.4% (17/386)   | 2.7% (88/3,291)     | 0.0% (0/53)         | 0.0% (0/30)     | 0.0% (0/33)      | 0.0% (0/39)   | 0.6% (1/170)    |
| Antimycobacterials         | Rifampicin (RIF, J04AH02)                  | 0.0% (0/39)       | 0.0% (0/37)         | 4.5% (300/6,709)    | 5.0% (28/562)   | 0.5% (25/4,549)     | 0.0% (0/82)         | 2.8% (1/36)     | 0.0% (0/39)      | 0.0% (0/34)   | 0.4% (1/229)    |
|                            | Amoxicillin (AMX, J01CA04)                 | 64.8% (107/165)   | 95.4% (3,026/3,173) | 95.4% (3,026/3,173) | 93.6% (280/299) | 92.8% (1,065/1,148) | 11.9% (5/42)        | 85.3% (29/34)   | 0.0% (0/37)      | 0.0% (0/34)   | 73.5% (36/49)   |
| Beta-lactams / penicillins | Amoxicillin/clavulanic acid (AMC, J01CR02) | 16.6% (107/643)   | 72.9% (3,026/4,153) | 72.9% (3,026/4,153) | 77.8% (280/360) | 46.1% (1,064/2,310) | 11.9% (5/42)        | 85.3% (29/34)   | 0.0% (0/37)      | 0.0% (0/34)   | 30.5% (36/118)  |
|                            | Ampicillin (AMP, J01CA01)                  | 64.8% (107/165)   | 95.4% (3,026/3,173) | 95.4% (3,026/3,173) | 93.6% (280/299) | 92.8% (1,065/1,148) | 11.9% (5/42)        | 85.3% (29/34)   | 0.0% (0/37)      | 0.0% (0/34)   | 73.5% (36/49)   |
|                            | Benzylpenicillin (PEN, J01CE01)            | 78.9% (870/1,102) | 97.1% (34/35)       | 92.5% (5,927/6,411) | 89.2% (486/545) | 88.2% (3,714/4,209) | 46.1% (1,064/2,310) | 57.5% (42/73)   | 25.0% (8/32)     | 0.0% (0/34)   | 77.8% (161/207) |
|                            | Flucloxacillin (FLC, J01CF05)              | 11.4% (140/1,227) | 55.3% (21/38)       | 61.8% (4,144/6,709) | 72.0% (406/564) | 37.2% (1,684/4,529) | 7.3% (6/82)         | 7.3% (6/82)     | 13.2% (5/38)     | 0.0% (0/34)   | 20.1% (46/229)  |
|                            | Oxacillin (OXA, J01CF04)                   | 13.4% (160/1,197) | 64.7% (22/34)       | 61.9% (4,135/6,679) | 72.1% (403/559) | 38.6% (1,692/4,386) | 7.3% (6/82)         | 7.3% (6/82)     | 16.2% (6/37)     | 0.0% (0/34)   | 20.5% (47/229)  |
|                            | Imipenem (IPM, J01DH15)                    | 21.8% (67/307)    | 73.4% (1,825/2,486) | 73.4% (1,825/2,486) | 79.8% (197/247) | 54.3% (506/932)     | 54.3% (506/932)     | 8.8% (5/57)     | 40.0% (14/35)    | 0.0% (0/34)   | 28.6% (16/56)   |
| Cephalosporins (1st gen.)  | Meropenem (MEM, J01DH02)                   | 21.8% (67/307)    | 73.4% (1,825/2,485) | 73.4% (1,825/2,485) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |
|                            | Cefazolin (CZO, J01DB04)                   | 21.8% (67/307)    | 73.4% (1,823/2,484) | 73.4% (1,823/2,484) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |
|                            | Cefosixin (FOX, J01DC01)                   | 21.8% (67/307)    | 73.4% (1,823/2,484) | 73.4% (1,823/2,484) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |
|                            | Cefturoxime (CXM, J01DC02)                 | 12.5% (104/831)   | 56.7% (17/30)       | 63.9% (3,025/4,733) | 70.2% (280/399) | 40.6% (1,063/2,621) | 40.6% (1,063/2,621) | 8.8% (5/57)     | 40.0% (14/35)    | 0.0% (0/34)   | 23.5% (36/153)  |
| Cephalosporins (3rd gen.)  | Cefotaxime (CTX, J01DD01)                  | 21.8% (67/307)    | 73.4% (1,824/2,485) | 73.4% (1,824/2,485) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |
|                            | Ceftaxone (CRO, J01DD04)                   | 21.8% (67/307)    | 73.4% (1,823/2,484) | 73.4% (1,823/2,484) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |
| Cephalosporins (4th gen.)  | Cefepime (FEP, J01DE01)                    | 21.8% (67/307)    | 73.4% (1,823/2,484) | 73.4% (1,823/2,484) | 79.8% (197/247) | 54.3% (506/932)     | 8.8% (5/57)         | 40.0% (14/35)   | 0.0% (0/34)      | 28.6% (16/56) | 28.6% (16/56)   |